

**FDA APPROVED**  
In individuals 16 years of age and older, the FDA-approved COMIRNATY® (COVID-19 Vaccine, mRNA) and the EUA-authorized Pfizer-BioNTech COVID-19 Vaccine have the same formulation and can be used interchangeably to provide the COVID-19 vaccination series.<sup>1</sup>

**AUTHORIZED FOR EMERGENCY USE**  
The emergency use of the authorized vaccine Pfizer-BioNTech COVID-19 Vaccine, under an Emergency Use Authorization (EUA), is for individuals 16 years of age and older; and the emergency use of this product is for individuals 12 years of age and older, and the emergency use of the medical product under Section 564(b)(1) of the FD&C Act unless the declaration is terminated or authorization revoked sooner.<sup>2</sup>

**Indication**

COMIRNATY® is a vaccine indicated for active immunization to prevent coronavirus disease 2019 (COVID-19) caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in individuals 16 years of age and older.

**Authorized Use**

COMIRNATY® (COVID-19 Vaccine, mRNA) is also authorized for emergency use to provide:

- a two-dose primary series in individuals 12 through 15 years
- a third primary series dose in individuals 12 years of age and older who have been determined to have certain kinds of immunocompromise
- a single booster dose in individuals:
  - 65 years of age and older

\*The licensed COMIRNATY® vaccine has the same formulation as the authorized vaccine Pfizer-BioNTech COVID-19 Vaccine, and the products can be used interchangeably to provide the vaccination series without presenting any safety or effectiveness concerns. The products are legally distinct, with certain differences that do not impact safety or effectiveness.

**Quick Access to Important Information**

- [Safety Information & Adverse Event Reporting](#)
- [Full Prescribing Information \(FDA Approved - 16 Years & Up\)](#)
- [EUA Fact Sheet for Vaccination Providers \(12 Years & Up\)](#)
- [EUA Fact Sheet for Recipients and Caregivers \(12 Years & Up\)](#)
- [Product Storage & Dry Ice](#)
- [Website for Recipients and Caregivers](#)

Tray remains consistent on every page of website

**Indication & Authorized Use**

The FDA-approved COMIRNATY® (COVID-19 Vaccine, mRNA) and the EUA-authorized Pfizer-BioNTech COVID-19 Vaccine have the same formulation and can be used interchangeably to provide the COVID-19 vaccination series.

**Indication**

COMIRNATY® is a vaccine indicated for active immunization to prevent coronavirus disease 2019 (COVID-19) caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in individuals 16 years of age and older.

**Authorized Use**

COMIRNATY® (COVID-19 Vaccine, mRNA) is also authorized for emergency use to provide:

**Important Safety Information**

Known history of a severe allergic reaction (eg, anaphylaxis) to any component of COMIRNATY® is a contraindication.

Appropriate medical treatment used to manage immediate allergic reactions must be immediately available in the event an acute anaphylactic reaction occurs following administration of the vaccine. Monitor vaccine recipients for the occurrence of immediate adverse reactions according to the Centers for Disease Control and Prevention guidelines (<https://www.cdc.gov/vaccines/covid-19/clinical-considerations/managing-anaphylaxis.html>).

Postmarketing data demonstrate increased risks of myocarditis and pericarditis, particularly within 7 days following the second dose.

Syncope (fainting) may occur in association with administration of injectable vaccines, including this vaccine. Procedures should be in place to avoid injury from fainting.

Immunocompromised persons, including individuals receiving immunosuppressant therapy, may have a diminished immune response to the vaccine.

The vaccine may not protect all vaccine recipients.

In clinical studies of participants 16 through 55 years of age, the most commonly reported adverse reactions (≥10%) were pain at the injection site (88.6%), fatigue (70.1%), headache (64.9%), muscle pain (45.5%), chills (41.5%), joint pain (27.5%), fever (17.8%), and injection site swelling (10.6%).

In clinical studies of participants 56 years of age and older, the most commonly reported adverse reactions (≥10%) were pain at the injection site (78.2%), fatigue (56.9%), headache (45.9%), muscle pain (32.5%), chills (24.8%), joint pain (21.5%), injection site swelling (11.8%), fever (11.5%), and injection site redness (10.4%).

In a clinical study evaluating administration of a booster dose in participants 18 through 55 years of age, the most commonly reported adverse reactions (≥10%) following the dose were pain at the injection



FDA APPROVED  
In individuals 16 years of age and older, the Pfizer-BioNTech COVID-19 Vaccine, mRNA (COMIRNATY®) (COVID-19 Vaccine, mRNA) is authorized for emergency use of this product to prevent COVID-19 in individuals 16 years of age and older; and the emergency use of this product to prevent COVID-19 in individuals 12 years of age and older to provide:

- a single booster dose in individuals:
  - 65 years of age and older
  - 18 through 64 years of age at high risk of severe COVID-19
  - 18 through 64 years of age whose frequent institutional or occupational exposure to SARS-CoV-2 puts them at high risk of serious complications of COVID-19 including severe COVID-19

The Pfizer-BioNTech COVID-19 Vaccine has received Emergency Use Authorization (EUA) from FDA to prevent COVID-19 in individuals 12 years of age and older to provide:

- a two-dose primary series in individuals 12 years of age and older
- a third primary series dose in individuals 12 years of age and older who have been determined to have certain kinds of immunocompromise
- a single booster dose in individuals:
  - 65 years of age and older
  - 18 through 64 years of age at high risk of severe COVID-19
  - 18 through 64 years of age whose frequent institutional or occupational exposure to SARS-CoV-2 puts them at high risk of serious complications of COVID-19 including severe COVID-19

AUTHORIZED FOR EMERGENCY USE  
The emergency use of the Pfizer-BioNTech COVID-19 Vaccine, mRNA (COMIRNATY®) (COVID-19 Vaccine, mRNA) is authorized for emergency use of this product to prevent COVID-19 in individuals 16 years of age and older; and the emergency use of this product to prevent COVID-19 in individuals 12 years of age and older to provide:

authorization of emergency use of the medical product under Section 564(b)(1) of the FD&C Act unless the declaration is terminated or authorization revoked sooner. <sup>2</sup>

<sup>1</sup>The licensed COMIRNATY® vaccine has the same formulation as the authorized vaccine Pfizer-BioNTech COVID-19 Vaccine, and the products can be used interchangeably to provide the vaccination series without presenting any safety or effectiveness concerns. The products are legally distinct, with certain differences that do not impact safety or effectiveness.

prevent COVID-19 caused by

FDA, under an Emergency Use Authorization (EUA) for individuals 12 years of age and older; and the emergency use of this product to prevent COVID-19 in individuals 12 years of age and older to provide:

most justifying the

## Quick Access to Important Information

 [Safety Information & Adverse Event Reporting](#)

 [Full Prescribing Information \(FDA Approved - 16 Years & Up\)](#)

 [EUA Fact Sheet for Vaccination Providers \(12 Years & Up\)](#)

 [EUA Fact Sheet for Recipients and Caregivers \(12 Years & Up\)](#)

 [Product Storage & Dry Ice](#)

 [Website for Recipients and Caregivers](#)

### Indication & Authorized Use

The FDA-approved COMIRNATY® (COVID-19 Vaccine, mRNA) and the EUA-authorized Pfizer-BioNTech COVID-19 Vaccine have the same formulation and can be used interchangeably to provide the COVID-19 vaccination series.

#### Indication

COMIRNATY® is a vaccine indicated for active immunization to prevent coronavirus disease 2019 (COVID-19) caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in individuals 16 years of age and older.

#### Authorized Use

COMIRNATY® (COVID-19 Vaccine, mRNA) is also authorized for emergency use to provide:

### Important Safety Information

Known history of a severe allergic reaction (eg, anaphylaxis) to any component of COMIRNATY® is a contraindication.

Appropriate medical treatment used to manage immediate allergic reactions must be immediately available in the event an acute anaphylactic reaction occurs following administration of the vaccine. Monitor vaccine recipients for the occurrence of immediate adverse reactions according to the Centers for Disease Control and Prevention guidelines (<https://www.cdc.gov/vaccines/covid-19/clinical-considerations/managing-anaphylaxis.html>).

Postmarketing data demonstrate increased risks of myocarditis and pericarditis, particularly within 7 days following the second dose.

Syncope (fainting) may occur in association with administration of injectable vaccines, including this vaccine. Procedures should be in place to avoid injury from fainting.

Immunocompromised persons, including individuals receiving immunosuppressant therapy, may have a diminished immune response to the vaccine.

The vaccine may not protect all vaccine recipients.

In clinical studies of participants 16 through 55 years of age, the most commonly reported adverse reactions (≥10%) were pain at the injection site (88.6%), fatigue (70.1%), headache (64.9%), muscle pain (45.5%), chills (41.5%), joint pain (27.5%), fever (17.8%), and injection site swelling (10.6%).

In clinical studies of participants 56 years of age and older, the most commonly reported adverse reactions (≥10%) were pain at the injection site (78.2%), fatigue (56.9%), headache (45.9%), muscle pain (32.5%), chills (24.8%), joint pain (21.5%), injection site swelling (11.8%), fever (11.5%), and injection site redness (10.4%).

In a clinical study evaluating administration of a booster dose in participants 18 through 55 years of age, the most commonly reported adverse reactions (≥10%) following the dose were pain at the injection



# FDA APPROVED FOR INDIVIDUALS >16 YEARS OF AGE<sup>1</sup>

# Quick Access to Important

## Indication & Authorized Use

The FDA-approved COMIRNATY® (COVID-19 Vaccine, mRNA) and the EUA-authorized Pfizer-BioNTech COVID-19 Vaccine have the same formulation and can be used interchangeably to provide the COVID-19 vaccination series.

### Indication

COMIRNATY® is a vaccine indicated for active immunization to prevent coronavirus disease 2019 (COVID-19) caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) in individuals 16 years of age and older.

### Authorized Use

COMIRNATY® (COVID-19 Vaccine, mRNA) is also authorized for emergency use to provide:

- a two-dose primary series in individuals 12 through 15 years
- a third primary series dose in individuals 12 years of age and older who have been determined to have certain kinds of immunocompromise
- a single booster dose in individuals:
  - 65 years of age and older
  - 18 through 64 years of age at high risk of severe COVID-19
  - 18 through 64 years of age whose frequent institutional or occupational exposure to SARS-CoV-2 puts them at high risk of serious complications of COVID-19 including severe COVID-19

The Pfizer-BioNTech COVID-19 Vaccine has received Emergency Use Authorization (EUA) from FDA to prevent COVID-19 in individuals 12 years of age and older to provide:

- a two-dose primary series in individuals 12 years of age and older
- a third primary series dose in individuals 12 years of age and older who have been determined to have certain kinds of immunocompromise
- a single booster dose in individuals:
  - 65 years of age and older
  - 18 through 64 years of age at high risk of severe COVID-19
  - 18 through 64 years of age whose frequent institutional or occupational exposure to SARS-CoV-2 puts them at high risk of serious complications of COVID-19 including severe COVID-19

## Important Safety Information

Known history of a severe allergic reaction (eg, anaphylaxis) to any component of COMIRNATY® is a contraindication.

Appropriate medical treatment used to manage immediate allergic reactions must be immediately available in the event an acute anaphylactic reaction occurs following administration of the vaccine.

Monitor vaccine recipients for the occurrence of immediate adverse reactions according to the Centers for Disease Control and Prevention guidelines (<https://www.cdc.gov/vaccines/covid-19/clinical-considerations/managing-anaphylaxis.html>).

Postmarketing data demonstrate increased risks of myocarditis and pericarditis, particularly within 7 days following the second dose.

Syncope (fainting) may occur in association with administration of injectable vaccines, including this vaccine. Procedures should be in place to avoid injury from fainting.

Immunocompromised persons, including individuals receiving immunosuppressant therapy, may have a diminished immune response to the vaccine.

The vaccine may not protect all vaccine recipients.

In clinical studies of participants 16 through 55 years of age, the most commonly reported adverse reactions (≥10%) were pain at the injection site (88.6%), fatigue (70.1%), headache (64.9%), muscle pain (45.5%), chills (41.5%), joint pain (27.5%), fever (17.8%), and injection site swelling (10.6%).

In clinical studies of participants 56 years of age and older, the most commonly reported adverse reactions (≥10%) were pain at the injection site (78.2%), fatigue (56.9%), headache (45.9%), muscle pain (32.5%), chills (24.8%), joint pain (21.5%), injection site swelling (11.8%), fever (11.5%), and injection site redness (10.4%).

In a clinical study evaluating administration of a booster dose in participants 18 through 55 years of age, the most commonly reported adverse reactions (≥10%) following the dose were pain at the injection site (83.0%), fatigue (63.7%), headache (48.4%), muscle pain (39.1%), chills (29.1%), and joint pain (25.3%).

In a clinical study of adolescents 12 through 15 years of age, the most commonly reported adverse reactions included pain at the injection site (90.5%), fatigue (77.5%), headache (75.5%), chills (49.2%), muscle pain (42.2%), fever (24.3%), joint pain (20.2%), injection site swelling (9.2%), injection site redness (8.6%), lymphadenopathy (0.8%), and nausea (0.4%).

Before administration of the vaccine, please see full [Prescribing Information](#) (16+ years of age), [EUA Fact Sheet for Vaccination Providers](#) (12+ years of age), and [Recipients and Caregivers Fact Sheet](#) (12+ years of age).

Patients should always ask their healthcare providers for medical advice about adverse events. You are encouraged to report negative side effects of vaccines to the US Food and Drug Administration (FDA) and the Centers for Disease Control and Prevention (CDC). Visit <https://vaers.hhs.gov/reportevent.html> or call 1-800-822-7967.

Drives to: <http://labeling.pfizer.com/ShowLabeling.aspx?id=15623>

Drives to: <http://labeling.pfizer.com/ShowLabeling.aspx?id=14471&format=pdf>

Drives to: <http://labeling.pfizer.com/ShowLabeling.aspx?id=14472&format=pdf>



## FDA APPROVED FOR INDIVIDUALS $\geq$ 16 YEARS OF AGE<sup>1</sup>

In individuals 16 years of age and older, the **Pfizer-BioNTech COVID-19 Vaccine, also known as COMIRNATY<sup>®</sup> (COVID-19 Vaccine, mRNA)**,\* has been approved for active immunization to prevent COVID-19 caused by SARS-CoV-2.<sup>1</sup>

## AUTHORIZED FOR EMERGENCY USE<sup>2</sup>

The emergency use of the product has not been approved or licensed by FDA, but has been authorized by FDA, under an Emergency Use Authorization (EUA) to prevent Coronavirus Disease 2019 (COVID-19) for use in individuals 12 years of age and older; and the emergency use of this product is only authorized for the duration of the declaration that circumstances exist justifying the authorization of emergency use of the medical product under Section 564(b)(1) of the FD&C Act unless the declaration is terminated or authorization revoked sooner.<sup>2</sup>

\*The licensed COMIRNATY<sup>®</sup> vaccine has the same formulation as the authorized vaccine Pfizer-BioNTech COVID-19 Vaccine, and the products can be used interchangeably to provide the vaccination series without presenting any safety or effectiveness concerns. The products are legally distinct, with certain differences that do not impact safety or effectiveness.

## Quick Access to Important Information

[Safety Information & Adverse Event Reporting](#)[Full Prescribing Information  
\(FDA Approved - 16 Years & Up\)](#)[EUA Fact Sheet for Vaccination Providers  
\(12 Years & Up\)](#)[EUA Fact Sheet for Recipients and Caregivers  
\(12 Years & Up\)](#)[Product Storage & Dry Ice](#)[Website for Recipients and Caregivers](#)

### References

1. COMIRNATY<sup>®</sup> (COVID-19 Vaccine, mRNA). Prescribing Information. Pfizer and BioNTech; 2021.
2. Pfizer-BioNTech COVID-19 Vaccine. Emergency Use Authorization Fact Sheet for Healthcare Providers Administering Vaccine (Vaccination Providers). Pfizer and BioNTech; 2021.

For more information

Pfizer Customer Service  
Call [1-800-879-3477](tel:1-800-879-3477).

Medical Information  
Visit [PfizerMedicalInformation.com](https://www.pfizer.com/medicalinformation) or call [1-800-438-1985](tel:1-800-438-1985).

General Product Inquiries  
Call [1-877-829-2619](tel:1-877-829-2619).

Shipment Support  
US Trade Customer Service  
Call [1-800-666-7248](tel:1-800-666-7248).